


ABSTRACT**PHARMACEUTICAL COMPOUNDS**

The invention provides a compound of the formula (I) for use in the prophylaxis or treatment of a disease state or condition mediated by a cyclin dependent kinase:



wherein

A is a group R^2 or CH_2-R^2 where R^2 is a carbocyclic or heterocyclic group having from 3 to 12 ring members;

B is a bond or an acyclic linker group having a linking chain length of up to 3 atoms selected from C, N, S and O;

R^1 is hydrogen or a group selected from SO_2R^b , $SO_2NR^7R^8$, $CONR^7R^8$, NR^7R^9 and carbocyclic and heterocyclic groups having from 3 to 7 ring members;

R^3 , R^4 , R^5 and R^6 are the same or different and are each selected from hydrogen, halogen, hydroxy, trifluoromethyl, cyano, nitro, carboxy, amino, carbocyclic and heterocyclic groups having from 3 to 12 ring members; a group R^a-R^b wherein R^a is a bond, O, CO, $X^1C(X^2)$, $C(X^2)X^1$, $X^1C(X^2)X^1$, S, SO, SO_2 , NR^c , SO_2NR^c or NR^cSO_2 ; and R^b is selected from hydrogen, carbocyclic and heterocyclic groups having from 3 to 12 ring members, and a C_{1-8} hydrocarbyl group optionally substituted by one or more substituents selected from hydroxy, oxo, halogen, cyano, nitro, amino, mono- or di- C_{1-4} hydrocarbylamino, carbocyclic and heterocyclic groups having from 3 to 12 ring members and wherein one or more carbon atoms of the C_{1-8} hydrocarbyl group may optionally be replaced by O, S, SO, SO_2 , NR^c , $X^1C(X^2)$, $C(X^2)X^1$ or $X^1C(X^2)X^1$;

R^c is hydrogen or C_{1-4} hydrocarbyl;

X^1 is O, S or NR^c and X^2 is =O, =S or = NR^c ;

- R^7 is selected from hydrogen and a C_{1-8} hydrocarbyl group optionally substituted by one or more substituents selected from hydroxy, oxo, halogen, cyano, nitro, amino, mono- or di- C_{1-4} hydrocarbylamino, carbocyclic and
- 5 heterocyclic groups having from 3 to 12 ring members and wherein one or more carbon atoms of the C_{1-8} hydrocarbyl group may optionally be replaced by O, S, SO, SO_2 , NR^c , $X^1C(X^2)$, $C(X^2)X^1$ or $X^1C(X^2)X^1$;
- R^8 is selected from R^7 and carbocyclic and heterocyclic groups having from 3 to 12 ring members;
- 10 R^9 is selected from R^8 , COR^8 and SO_2R^8 ;
- or NR^7R^8 or NR^7R^9 may each form a heterocyclic group having from 5 to 12 ring members;
- but excluding the compounds N-[(morpholin-4-yl)phenyl-1H-indazole-3-carboxamide and N-[4-(acetaminosulphonyl)phenyl-1H-indazole-3-carboxamide.